













































Common Panhandle Wildflowers

 <p><i>Conoclinium coelestinum</i> Blue mistflower Blue mistflower is a sure sign that autumn has arrived or is certainly not far away. Spreading rapidly by underground runners, this aggressive herbaceous perennial forms large colonies of 1'-3' tall plants. The numerous compact heads of deep blue or purple flowers add significant charm to October roadsides. Fall</p>	 <p><i>Symphyotrichum adnatum</i> Scaleleaf aster This is one of several blue-flowered asters found along Panhandle roadsides. It is a low-growing species with tiny, awl-shaped leaves that point upward and tightly hug the stem. Chapman's aster (<i>Symphyotrichum chapmani</i>) has similar flowers but grows much taller, with a slender arching stem that is nearly devoid of leaves. Spring, Summer, Fall</p>	 <p><i>Amsonia ciliata</i> Fringed bluestar Two species of bluestar occur in the Panhandle, one in wet woodlands, and the other along the backslope of dry, sandy roadsides. The stem is about 2' tall with very narrow leaves. It is topped by a showy cluster of pale blue to bluish-white star-shaped flowers. Spring</p>	 <p><i>Salvia lyrata</i> Lyreleaf sage The lavender-blue flowers of Lyreleaf sage often dominate early spring roadsides. Plants average 1'-2' tall with a single upright stalk and a whorl-like cluster of ground-hugging leaves. Spring</p>	 <p><i>Carphephorus odoratissimus</i> Vanillaleaf This single-stemmed, fall-blooming herb ranges from 1'-4' tall and has numerous densely packed heads of tiny purple flowers. It typically occurs in large, conspicuous colonies that are especially notable when flowers are present. It is known as Vanillaleaf because of the scent of its leaves. Fall</p>
 <p><i>Liatris</i> Blazing star About a dozen species of blazing stars occur in the central Panhandle, all of which have erect, spike-like inflorescences. Species vary from 2'-6' tall and usually occur in showy colonies along moist or dry roadsides. The individual flowers are small, 5-lobed, and star-shaped, which explains the common name. Pictured: Chapman's blazing star (<i>Liatris chapmani</i>). Summer, Fall</p>	 <p><i>Cirsium</i> Thistle The several thistles that occur along Florida's roadsides are among our most distinctive wildflowers. Although their flower structure is similar, flower color can range from yellow to pinkish or purplish. Purple thistle (<i>Cirsium horridulum</i>), pictured, is characterized by a treacherous nest of prickly, leaflike bracts under the flower head. Spring, Summer</p>	 <p><i>Vernonia</i> Ironweed Ironweeds are tall, wispy plants with striking purple flowers and an upright form. Plants average 3'-5' tall, usually with several upright stems. Some prefer dry sites, others moist roadsides. Fall</p>	 <p><i>Lupinus villosus</i> Lady lupine Lady lupine is one of the more common of our native lupines. The dense cluster of hairy, blue-gray leaves and erect 1'-2' spikes of lavender to pinkish flowers make a conspicuous display along dry, sandy roadsides. Spring</p>	 <p><i>Monarda punctata</i> Dotted horsemint Dotted horsemint is worth stopping for. The large purplish-pink bracts and spotted yellow flowers are exquisite when seen close up, and the minty aroma only adds to the sensory appeal. Handling any part of the plant can impart a pleasing fragrance to the skin. Fall</p>
 <p><i>Agalinis</i> False foxglove The attractive lavender or purple flowers of our several false foxgloves are harbingers of autumn. Most begin to flower in late summer but are at their best in September and October. About a dozen species occur in our region, many of which are difficult to distinguish from each other. Summer, Fall</p>	 <p><i>Rhexia</i> Meadowbeauty As their name suggests, meadowbeauties prefer treeless savannas, open bogs and sunny roadsides. There are about a dozen species, most of which have pink or purple flowers with large yellow hooklike anthers. They are one of our most abundant groups of summer-blooming wildflowers. Pictured: Nash's meadowbeauty (<i>Rhexia nashii</i>). Summer</p>	 <p><i>Sabatia</i> Rosegentian Rosegentians are summer delights. About a dozen species occur in the Panhandle, some with white flowers, others with pink flowers. Roadside species usually have five or 10 petals. Bartram's rosegentian (<i>Sabatia decandra</i>), pictured, is bright pink with 10 petals. Summer</p>	 <p><i>Coreopsis nudata</i> Pink Coreopsis Our only pink-flowered Coreopsis is found mostly in the Panhandle. Plants can grow to 5' tall and often occur in large showy colonies, with numerous flowerheads adding a conspicuous splash of color to wet ditches. Pink Coreopsis is common along several of this brochure's recommended routes. Spring</p>	 <p><i>Calopogon barbatus</i> Grasspink orchid Several species of grasspink orchids occur along the routes recommended here. All have "upside-down" flowers, with the lip petal pointing upward rather than downward. The patch of showy hairs at the top of the lip resembles stamens and are an adaptation designed to attract bees and other pollinators. Pictured: Bearded grasspink orchid (<i>Calopogon barbatus</i>). Spring</p>
 <p><i>Ipomoea cordatotriloba</i> Tievine Like the showy blossoms of most morning glories, Tievine flowers fade quickly as the day progresses. Its flowers are funnel-shaped, less than 2" wide, and range in color from pink to rose-purple. It is conspicuous as it scrambles over roadside vegetation in late summer and autumn. Summer, Fall</p>	 <p><i>Stenanthium densum</i> Crowpoison, Osceola's plume Natural and prescribed fire is a well-known wildflower catalyst. This is certainly true of Crowpoison, which sends up its slender 3' stems and cone-shaped inflorescences within days or weeks following a growing-season burn. The small individual flowers start out white, but turn purplish as they age. Spring</p>	 <p><i>Zephyranthes</i> Rainily Rainilies often flower immediately following a downpour. Two similar species occur in our region. Both have shiny green grasslike leaves and white trumpet-shaped flowers. One species grows along the margins of moist roadside ditches, the other mostly in shady hardwood forests. Spring</p>	 <p><i>Eupatorium</i> Thoroughwort, Fennel At least half of the nearly 20 thoroughworts that grace Florida's roadsides occur in the Panhandle. All have tiny white flowers borne in congested heads at the top of a branched inflorescence. Pictured: Common boneset (<i>Eupatorium perfoliatum</i>). Summer, Fall</p>	 <p><i>Baptisia alba</i> White wild indigo This large, shrubby bean stands 3'-4' tall. In early to mid-spring it produces numerous tall, showy racemes of creamy white pea flowers that are unmistakable. It prefers moist, open roadsides in full sun. Mid-Spring</p>
 <p><i>Erigeron</i> Daisy fleabane The daisy fleabanes are among the most common roadside wildflowers. They are easily recognized by their daisylike flowerheads with a dark center surrounded by numerous small, laterally spreading rays. At least four species occur in the Panhandle. Early whitetop fleabane (<i>Erigeron vernus</i>), pictured, blooms in spring. Spring, Summer</p>	 <p><i>Eriocaulon decangulare</i> Tenangle pipewort The erect 1'-3' tall arrow-straight stem, topped by a buttonlike cluster of creamy-white flowers, suggests the decorative pin used to hold a lady's hat to her head. These clump-forming perennials are denizens of wet roadside ditches and often occur in large, conspicuous colonies. Spring, Summer, Fall</p>	 <p><i>Eurybia eryngiifolia</i> Thistleleaf aster The disheveled appearance of the thin creamy-white rays surrounding a dark-brown center sets this species apart from other asters. Its Florida range is restricted to the several counties of the central Panhandle, and it is common along roadsides in early summer, especially in the Apalachicola National Forest. Spring, Summer</p>	 <p><i>Rhynchospora latifolia</i> Whitetop sedge Not many sedges make the list of showy bloomers. <i>Rhynchospora latifolia</i>, pictured, is one of two species of Whitetop sedge that are the exceptions. The white structures below the flowers are actually leaflike bracts, but their effect is that of flower petals. Spring, Summer</p>	 <p><i>Lachnanthes caroliniana</i> Redroot Redroot makes its appearance in late May, with its peak flowering in mid- to late June. Its 2' stem is crowned by a white and yellow inflorescence and is very showy. Large populations of several hundred plants often occupy wet roadside ditches and are easily spotted, even at highway speed. Summer</p>
 <p><i>Lophiola aurea</i> Golden crest This wetland-loving herb stands 1'-3' tall and prefers savannas and wet ditches. The whitish, hairy stem supports a branching inflorescence. The scientific epithet <i>aurea</i> means golden, a fitting description of the flower color. Spring, Summer, Fall</p>	 <p><i>Pyrrohapappus carolinianus</i> Carolina desert chicory This yellow-flowered member of the Aster family begins blooming as early as February and continues throughout the spring, even into mid-summer. Its slender stem is about 2' tall and is terminated by a dense head consisting only of soft yellow ray flowers. Unlike the Black-eyed Susan and most other species in this family, the flower head lacks a central disk. Spring, Summer</p>	 <p><i>Hypericum</i> St. John's wort More than 20 species of St. John's wort occur in the Panhandle, many along roadsides and woodland margins. All have yellow 4- or 5-petaled flowers with a brushlike mass of yellow stamens. Peckbank St. John's wort (<i>Hypericum fasciculatum</i>), pictured, flowers in the spring. Spring, Summer, Fall</p>	 <p><i>Packera glabella</i> Butterweed Butterweed is one of our earliest blooming native wildflowers, sometimes flowering as early as January. It stands about 2' tall and prefers low roadsides and wet ditches. The golden flowers and deeply divided leaves are distinctive. Spring</p>	 <p><i>Coreopsis</i> Coreopsis, Tickseed Coreopsis is Florida's state wildflower. Sixteen species occur in the state. Most, like this Lanceleaf Coreopsis (<i>Coreopsis lanceolata</i>), have yellow disk flowers with ray flowers notched at the tips. Lanceleaf Coreopsis is one of Florida's more common, showy spring wildflowers. Spring, Summer, Fall</p>
 <p><i>Balduina</i> Honeycombhead Honeycombhead usually occurs in large showy populations, with thousands of plants stretching across the wet savannas of the Apalachicola National Forest. Two species occur in the Panhandle, including one that prefers dry, sandy roadsides. Fall</p>	 <p><i>Helianthus heterophyllus</i> Sunflower This sunflower is among the showiest of Florida's fall wildflowers, decorating moist road shoulders and wet roadside ditches. The state is home to other <i>Helianthus</i> species, most of which have yellow petals circling a brown or yellow center. The only exception is the Rayless sunflower (pictured below). Fall</p>	 <p><i>Rudbeckia mohrii</i> Mohr's coneflower Several of our roadside <i>Rudbeckias</i> are immediately recognizable from their resemblance to garden plants of the same name. Most have a central conelike head surrounded by numerous yellow rays. Mohr's coneflower is one of our showiest examples. Summer</p>	 <p><i>Berlandiera pumila</i> Soft greeneyes Soft greeneyes takes its name from its flowerheads, which have a round green disk at the center of eight bright yellow rays. The stem is 8"-32" tall with conspicuous 4" leaves. It is most often seen on dry, open, sandy roadsides. Disk flowers have a chocolate aroma when open. Spring, Summer</p>	 <p><i>Aletris lutea</i> Yellow colicroot Yellow colicroot can dominate moist to wet roadsides during its mid- to late spring flowering period. The stiffly erect, 3' stem arises from a flattened whorl of ground-hugging leaves and is terminated by a spike-like inflorescence of bright yellow, tubular flowers. Spring</p>
 <p><i>Helenium</i> Sneezeweed Several species of sneezeweed are found along Florida roadsides. Savanna sneezeweed (<i>Helenium vernale</i>), pictured, is about 3' tall and blooms in mid-spring. It is typical of the group. At least one species flowers in summer. Spring, Summer</p>	 <p><i>Solidago</i> Goldenrod The narrow spike-like inflorescences of goldenrods give them their common name. They are among our most recognized native wildflowers and usually bloom in the fall. Species vary from 2'-4' tall and prefer open roadsides in full sun. Fall</p>	 <p><i>Sarracenia</i> Pitcherplants Northwest Florida is well known for its assortment of insectivorous pitcherplants, at least four species of which can be seen along our recommended roadsides. The trumpet-like leaves of the Yellow pitcherplant (<i>Sarracenia flava</i>), shown here, can be 2' tall. All can be seen along wet to moist roadsides, especially along SR 65 and the Apalachee Savannas Scenic Byway. Spring, Summer</p>	 <p><i>Platanthera</i> Fringed and Fringeless orchid Several of these upright, 1'-3' tall orchids can be found along moist or boggy roadsides, especially in the Apalachicola National Forest. Flower color ranges from bright white to yellow or orange. Blooming begins in June for the white-flowered forms, with a succession of oranges and yellows from mid-July into August. Pictured: Crested fringed orchid (<i>Platanthera cristata</i>). Spring, Summer</p>	 <p>Summer <i>Asclepias lanceolata</i> Lanceleaf milkweed Milkweeds are most easily recognized by their unique flowers and best known for their attractiveness to butterflies. Some are low-growing; others tall and conspicuous. Flower color ranges from white to lavender, green, yellow, or bright orange-red. Lanceleaf milkweed can be 3'-4' tall, which makes it a conspicuous component of wet roadside ditches. Spring, Summer</p>
 <p><i>Campsis radicans</i> Trumpet creeper The tubular orange-red flowers of Trumpet creeper make their first appearance in late spring but continue into summer. This is a high-climbing woody vine with lacy leaves. Look for it scaling trees and shrubs just off the road shoulder. Spring, Summer</p>	 <p><i>Helianthus radula</i> Rayless sunflower Rayless sunflower stands in stark contrast to Florida's other <i>Helianthus</i> species. Although its flowerhead lacks the bright yellow rays associated with most species, its dark brown disk more than makes up for the loss. Its sheer abundance in wet savannas and ditches ensures it won't be missed. Fall</p>	 <p><i>Rudbeckia graminifolia</i> Grassleaf coneflower The combination of short brick-red rays and large conelike disk sets Grassleaf coneflower apart. It can be abundant in savannas and on moist roadsides, where hundreds of the 2' stems can dot the landscape, each topped with a single flower head. Summer</p>	 <p><i>Trillium</i> Wakerobin You'll have to find a shady hardwood forest to see our native trilliums. The leaves of these long-lived perennials make their annual appearance in late winter and flower in early spring, well before the trees put on new leaves. Chattahoochee Nature Park, Torreya State Park and Jackson Mounds are good destinations to search for these interesting plants. Spring</p>	